

Local proof-theoretic foundations and proof-theoretic tameness in ordinary mathematics

Ulrich Kohlenbach

Department of Mathematics



TECHNISCHE
UNIVERSITÄT
DARMSTADT

ASL Logic Colloquium 2019, Prague, August 15, 2019

Retiring Presidential Address

Proof Mining since 2000 (abstract classes of spaces)

Since K. NFAO 2001: applications of proof mining in analysis with **abstract metric spaces** involved: **fixed point and ergodic theory, convex optimization, geodesic geometry, Cauchy problems, game theory etc.**

Proof Mining since 2000 (abstract classes of spaces)

Since K. NFAO 2001: applications of proof mining in analysis with **abstract metric spaces** involved: **fixed point and ergodic theory, convex optimization, geodesic geometry, Cauchy problems, game theory etc.**

Setting: X normed linear or geodesic space, $C \subseteq X$ some (often convex) subset and $T : C \rightarrow C$ or $F : C \rightarrow \overline{\mathbb{R}}$ nonlinear maps.

Proof Mining since 2000 (abstract classes of spaces)

Since K. NFAO 2001: applications of proof mining in analysis with **abstract metric spaces** involved: **fixed point and ergodic theory, convex optimization, geodesic geometry, Cauchy problems, game theory etc.**

Setting: X normed linear or geodesic space, $C \subseteq X$ some (often convex) subset and $T : C \rightarrow C$ or $F : C \rightarrow \overline{\mathbb{R}}$ nonlinear maps.

Covers numerous fixed point, zero-finding, minimization or equilibrium problems with iterative procedures (x_n) s.t. e.g. in the case of fixed point problems one has

$$(1) \ d(x_n, Tx_n) \xrightarrow{n \rightarrow \infty} 0 \text{ or even}$$

$$(2) \ (x_n) \text{ strongly converges to the fixed point of } T.$$

For such situations, **special designed** (for **particular classes** of spaces X and mappings T) **logical metatheorems** (K. TAMS 2005, Gerhardy/K. TAMS 2008) have been designed which guarantee the extractability of explicit uniform bounds for $\forall \underline{x} \in \mathbb{N}, \mathbb{N}^{\mathbb{N}}, \mathbf{X}, \mathbf{X}^{\mathbf{X}}, \mathbf{X}^{\mathbb{N}} \dots \exists n \in \mathbb{N} \mathbf{A}(\underline{x}, n)$ -theorems.

For such situations, **special designed** (for **particular classes** of spaces X and mappings T) **logical metatheorems** (K. TAMS 2005, Gerhardy/K. TAMS 2008) have been designed which guarantee the extractability of explicit uniform bounds for $\forall \underline{x} \in \mathbb{N}, \mathbb{N}^{\mathbb{N}}, \mathbf{X}, \mathbf{X}^{\mathbf{X}}, \mathbf{X}^{\mathbb{N}} \dots \exists n \in \mathbb{N} \mathbf{A}(\underline{x}, n)$ -theorems. Due to **classical logic**: in general \mathbf{A} must be **existential** \mathbf{A}_{\exists} .

For such situations, **special designed** (for **particular classes** of spaces X and mappings T) **logical metatheorems** (K. TAMS 2005, Gerhardy/K. TAMS 2008) have been designed which guarantee the extractability of explicit uniform bounds for $\forall \underline{x} \in \mathbb{N}, \mathbb{N}^{\mathbb{N}}, \mathbf{X}, \mathbf{X}^{\mathbf{X}}, \mathbf{X}^{\mathbb{N}} \dots \exists n \in \mathbb{N} \mathbf{A}(\underline{x}, n)$ -theorems.

Due to **classical logic**: in general \mathbf{A} must be **existential** \mathbf{A}_{\exists} .

For **restricted use of LEM arbitrary \mathbf{A} !**

For such situations, **special designed** (for **particular classes** of spaces X and mappings T) **logical metatheorems** (K. TAMS 2005, Gerhardy/K. TAMS 2008) have been designed which guarantee the extractability of explicit uniform bounds for $\forall \underline{x} \in \mathbb{N}, \mathbb{N}^{\mathbb{N}}, \mathbf{X}, \mathbf{X}^{\mathbf{X}}, \mathbf{X}^{\mathbb{N}} \dots \exists n \in \mathbb{N} \mathbf{A}(\underline{x}, n)$ -theorems.

Due to **classical logic**: in general \mathbf{A} must be **existential** \mathbf{A}_{\exists} .

For **restricted use of LEM arbitrary \mathbf{A} !**

The **logical metatheorems** guarantee the extractability of effective bounds on ' \exists ' **independent** from parameters in

For such situations, **special designed** (for **particular classes** of spaces X and mappings T) **logical metatheorems** (K. TAMS 2005, Gerhardy/K. TAMS 2008) have been designed which guarantee the extractability of explicit uniform bounds for $\forall \underline{x} \in \mathbb{N}, \mathbb{N}^{\mathbb{N}}, X, X^X, X^{\mathbb{N}} \dots \exists n \in \mathbb{N} A(\underline{x}, n)$ -theorems.

Due to **classical logic**: in general A must be **existential** A_{\exists} .

For **restricted use of LEM arbitrary** A !

The **logical metatheorems** guarantee the extractability of effective bounds on ‘ \exists ’ **independent** from parameters in

- **compact** metric spaces (**if separability is used**) and

For such situations, **special designed** (for **particular classes** of spaces X and mappings T) **logical metatheorems** (K. TAMS 2005, Gerhardy/K. TAMS 2008) have been designed which guarantee the extractability of explicit uniform bounds for $\forall \underline{x} \in \mathbb{N}, \mathbb{N}^{\mathbb{N}}, X, X^X, X^{\mathbb{N}} \dots \exists n \in \mathbb{N} A(\underline{x}, n)$ -theorems.

Due to **classical logic**: in general A must be **existential** A_{\exists} .

For **restricted use of LEM arbitrary A !**

The **logical metatheorems** guarantee the extractability of effective bounds on ' \exists ' **independent** from parameters in

- **compact** metric spaces (**if separability is used**) and
- bounded subsets of **abstract** metric structures X .

Formal systems for analysis with abstract metric spaces X

Types: (i) \mathbb{N}, X are types, (ii) with ρ, τ also $\rho \rightarrow \tau$ is a type.

Formal systems for analysis with abstract metric spaces X

Types: (i) \mathbb{N}, X are types, (ii) with ρ, τ also $\rho \rightarrow \tau$ is a type.

$\mathbf{PA}^{\omega, X}$ is the extension of Peano Arithmetic to all types.

$\mathcal{A}^{\omega, X} := \mathbf{PA}^{\omega, X} + \mathbf{Dependent\ choice\ in\ all\ types}.$

Formal systems for analysis with abstract metric spaces X

Types: (i) \mathbb{N}, X are types, (ii) with ρ, τ also $\rho \rightarrow \tau$ is a type.

$\mathbf{PA}^{\omega, X}$ is the extension of Peano Arithmetic to all types.

$\mathcal{A}^{\omega, X} := \mathbf{PA}^{\omega, X} + \text{Dependent choice in all types.}$

Implies **full comprehension** for numbers (2nd order arithmetic).

In practice: only **weak fragments** needed (after pre-processing).

Formal systems for analysis with abstract metric spaces X

Types: (i) \mathbb{N}, X are types, (ii) with ρ, τ also $\rho \rightarrow \tau$ is a type.

$\mathbf{PA}^{\omega, X}$ is the extension of Peano Arithmetic to all types.

$\mathcal{A}^{\omega, X} := \mathbf{PA}^{\omega, X} + \text{Dependent choice in all types.}$

Implies **full comprehension** for numbers (2nd order arithmetic).

In practice: only **weak fragments** needed (after pre-processing).

Equality is a **defined** notion: $x^X =_X y^X \equiv d_X(x, y) =_{\mathbb{R}} 0_{\mathbb{R}}$.

Formal systems for analysis with abstract metric spaces X

Types: (i) \mathbb{N}, X are types, (ii) with ρ, τ also $\rho \rightarrow \tau$ is a type.

$\mathbf{PA}^{\omega, X}$ is the extension of Peano Arithmetic to all types.

$\mathcal{A}^{\omega, X} := \mathbf{PA}^{\omega, X} + \text{Dependent choice in all types.}$

Implies **full comprehension** for numbers (2nd order arithmetic).

In practice: only **weak fragments** needed (after pre-processing).

Equality is a **defined** notion: $x^X =_X y^X \equiv d_X(x, y) =_{\mathbb{R}} 0_{\mathbb{R}}$.

In general only **rule**

$$\frac{s =_X t}{T(s) =_X T(t)}.$$

Formal systems for analysis with abstract metric spaces X

Types: (i) \mathbb{N}, X are types, (ii) with ρ, τ also $\rho \rightarrow \tau$ is a type.

$\mathbf{PA}^{\omega, X}$ is the extension of Peano Arithmetic to all types.

$\mathcal{A}^{\omega, X} := \mathbf{PA}^{\omega, X} + \text{Dependent choice in all types.}$

Implies **full comprehension** for numbers (2nd order arithmetic).

In practice: only **weak fragments** needed (after pre-processing).

Equality is a **defined** notion: $x^X =_X y^X \equiv d_X(x, y) =_{\mathbb{R}} 0_{\mathbb{R}}$.

In general only **rule**

$$\frac{s =_X t}{T(s) =_X T(t)}.$$

$\mathcal{A}^{\omega}[X, \|\cdot\| \dots]$ e.g. results by adding constants with axioms expressing that $(X, \|\cdot\|)$ is normed, uniformly convex, Hilbert.

Keeping track of uniform bounds: majorization

y, x functionals of types $\rho, \hat{\rho} := \rho[\mathbb{N}/X]$:

$$\begin{aligned} \mathbf{x}^{\mathbb{N}} \gtrsim_{\mathbb{N}} \mathbf{y}^{\mathbb{N}} &: \equiv \mathbf{x} \geq \mathbf{y} \\ \mathbf{x}^{\mathbb{N}} \gtrsim_{\mathbf{x}} \mathbf{y}^{\mathbf{x}} &: \equiv \mathbf{x} \geq \|\mathbf{y}\|. \end{aligned}$$

Keeping track of uniform bounds: majorization

y, x functionals of types $\rho, \hat{\rho} := \rho[\mathbb{N}/X]$:

$$\begin{aligned}x^{\mathbb{N}} \gtrsim_{\mathbb{N}} y^{\mathbb{N}} &: \equiv x \geq y \\x^{\mathbb{N}} \gtrsim_x y^x &: \equiv x \geq \|y\|.\end{aligned}$$

For **complex types** $\rho \rightarrow \tau$ this is extended in a **hereditarily**.

Keeping track of uniform bounds: majorization

y, x functionals of types $\rho, \hat{\rho} := \rho[\mathbb{N}/X]$:

$$\begin{aligned}x^{\mathbb{N}} \gtrsim_{\mathbb{N}} y^{\mathbb{N}} &: \equiv x \geq y \\x^{\mathbb{N}} \gtrsim_x y^x &: \equiv x \geq \|y\|.\end{aligned}$$

For **complex types** $\rho \rightarrow \tau$ this is extended in a **hereditarily**.

Example:

$$f^* \gtrsim_{x \rightarrow x} f \equiv \forall n \in \mathbb{N}, x \in X [n \geq \|x\| \rightarrow f^*(n) \geq \|f(x)\|].$$

Keeping track of uniform bounds: majorization

y, x functionals of types $\rho, \hat{\rho} := \rho[\mathbb{N}/X]$:

$$\begin{aligned}x^{\mathbb{N}} \gtrsim_{\mathbb{N}} y^{\mathbb{N}} &: \equiv x \geq y \\x^{\mathbb{N}} \gtrsim_x y^x &: \equiv x \geq \|y\|.\end{aligned}$$

For **complex types** $\rho \rightarrow \tau$ this is extended in a **hereditarily**.

Example:

$$f^* \gtrsim_{X \rightarrow X} f \equiv \forall n \in \mathbb{N}, x \in X [n \geq \|x\| \rightarrow f^*(n) \geq \|f(x)\|].$$

$$f : X \rightarrow X \text{ **nonexpansive**: } \|f(x) - f(y)\| \leq \|x - y\|.$$

Keeping track of uniform bounds: majorization

y, x functionals of types $\rho, \hat{\rho} := \rho[\mathbb{N}/X]$:

$$\begin{aligned}x^{\mathbb{N}} \gtrsim_{\mathbb{N}} y^{\mathbb{N}} &: \equiv x \geq y \\x^{\mathbb{N}} \gtrsim_X y^X &: \equiv x \geq \|y\|.\end{aligned}$$

For **complex types** $\rho \rightarrow \tau$ this is extended in a **hereditarily**.

Example:

$$f^* \gtrsim_{X \rightarrow X} f \equiv \forall n \in \mathbb{N}, x \in X [n \geq \|x\| \rightarrow f^*(n) \geq \|f(x)\|].$$

$$f : X \rightarrow X \text{ **nonexpansive**: } \|f(x) - f(y)\| \leq \|x - y\|.$$

Then for $\|f(0)\| \leq b$ and $f^*(n) := n + b$: $f^* \gtrsim_{X \rightarrow X} f$.

Keeping track of uniform bounds: majorization

y, x functionals of types $\rho, \hat{\rho} := \rho[\mathbb{N}/X]$:

$$\begin{aligned}x^{\mathbb{N}} \gtrsim_{\mathbb{N}} y^{\mathbb{N}} &: \equiv x \geq y \\x^{\mathbb{N}} \gtrsim_X y^X &: \equiv x \geq \|y\|.\end{aligned}$$

For **complex types** $\rho \rightarrow \tau$ this is extended in a **hereditarily**.

Example:

$$f^* \gtrsim_{X \rightarrow X} f \equiv \forall n \in \mathbb{N}, x \in X [n \geq \|x\| \rightarrow f^*(n) \geq \|f(x)\|].$$

$$f : X \rightarrow X \text{ **nonexpansive**: } \|f(x) - f(y)\| \leq \|x - y\|.$$

Then for $\|f(0)\| \leq b$ and $f^*(n) := n + b$: $f^* \gtrsim_{X \rightarrow X} f$.

In a **metric setting**: **reference point** $a \in X$: $x \geq d(a, y)$.

Special case of **general logical metatheorems** (T nonexpansive):

Corollary (Gerhardy/K., TAMS 2008)

If $\mathcal{A}^\omega[X, \|\cdot\|]$ proves (K represented compact metric space)

$\forall x \in \mathbb{N}^{\mathbb{N}} \forall y \in K \forall z \in X \forall T : X \rightarrow X \left(T \text{ n.e.} \rightarrow \exists v \in \mathbb{N} A_{\exists} \right),$

Special case of **general logical metatheorems** (T nonexpansive):

Corollary (Gerhardy/K., TAMS 2008)

If $\mathcal{A}^\omega[X, \|\cdot\|]$ proves (K represented compact metric space)

$$\forall x \in \mathbb{N}^\mathbb{N} \forall y \in K \forall z \in X \forall T : X \rightarrow X \left(T \text{ n.e.} \rightarrow \exists v \in \mathbb{N} A_\exists \right),$$

then one can extract a **computable functional** $\Phi : \mathbb{N}^\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$
s.t. in all normed spaces X it holds that

$$\forall x \in \mathbb{N}^\mathbb{N} \forall y \in K \forall z \in X \forall T : X \rightarrow X \forall b \in \mathbb{N} \\ \left(T \text{ n.e.} \wedge \|z\|, \|T(0)\| \leq b \rightarrow \exists v \leq \Phi(x, b) A_\exists \right).$$

Special case of **general logical metatheorems** (T nonexpansive):

Corollary (Gerhardy/K., TAMS 2008)

If $\mathcal{A}^\omega[X, \|\cdot\|]$ proves (K represented compact metric space)

$$\forall x \in \mathbb{N}^{\mathbb{N}} \forall y \in K \forall z \in X \forall T : X \rightarrow X \left(T \text{ n.e.} \rightarrow \exists v \in \mathbb{N} A_{\exists} \right),$$

then one can extract a **computable functional** $\Phi : \mathbb{N}^{\mathbb{N}} \times \mathbb{N} \rightarrow \mathbb{N}$
s.t. in all normed spaces X it holds that

$$\forall x \in \mathbb{N}^{\mathbb{N}} \forall y \in K \forall z \in X \forall T : X \rightarrow X \forall b \in \mathbb{N} \\ \left(T \text{ n.e.} \wedge \|z\|, \|T(0)\| \leq b \rightarrow \exists v \leq \Phi(x, b) A_{\exists} \right).$$

Similar for Hilbert spaces and uniformly convex spaces (then bound depends on modulus of convexity).

Special case of **general logical metatheorems** (T nonexpansive):

Corollary (Gerhardy/K., TAMS 2008)

If $\mathcal{A}^\omega[X, \|\cdot\|]$ proves (K represented compact metric space)

$$\forall x \in \mathbb{N}^{\mathbb{N}} \forall y \in K \forall z \in X \forall T : X \rightarrow X \left(T \text{ n.e.} \rightarrow \exists v \in \mathbb{N} A_{\exists} \right),$$

then one can extract a **computable functional** $\Phi : \mathbb{N}^{\mathbb{N}} \times \mathbb{N} \rightarrow \mathbb{N}$
s.t. in all normed spaces X it holds that

$$\forall x \in \mathbb{N}^{\mathbb{N}} \forall y \in K \forall z \in X \forall T : X \rightarrow X \forall b \in \mathbb{N} \\ \left(T \text{ n.e.} \wedge \|z\|, \|T(0)\| \leq b \rightarrow \exists v \leq \Phi(x, b) A_{\exists} \right).$$

Similar for Hilbert spaces and uniformly convex spaces (then bound depends on modulus of convexity). In metric setting: $d(z, Tz) \leq b$.

Special case of **general logical metatheorems** (T nonexpansive):

Corollary (Gerhardy/K., TAMS 2008)

If $\mathcal{A}^\omega[X, \|\cdot\|]$ proves (K represented compact metric space)

$$\forall x \in \mathbb{N}^{\mathbb{N}} \forall y \in K \forall z \in X \forall T : X \rightarrow X \left(T \text{ n.e.} \rightarrow \exists v \in \mathbb{N} A_{\exists} \right),$$

then one can extract a **computable functional** $\Phi : \mathbb{N}^{\mathbb{N}} \times \mathbb{N} \rightarrow \mathbb{N}$
s.t. in all normed spaces X it holds that

$$\forall x \in \mathbb{N}^{\mathbb{N}} \forall y \in K \forall z \in X \forall T : X \rightarrow X \forall b \in \mathbb{N} \\ \left(T \text{ n.e.} \wedge \|z\|, \|T(0)\| \leq b \rightarrow \exists v \leq \Phi(x, b) A_{\exists} \right).$$

Similar for Hilbert spaces and uniformly convex spaces (then bound depends on modulus of convexity). In metric setting: $d(z, Tz) \leq b$.

Method: Novel forms of Gödel's functional interpretation!

Applicability of Metatheorem

- Applied to asymptotic regularity statements $d(x_n, Tx_n) \rightarrow 0$, the corollary often gives full rates of convergence, e.g. because $(d(x_n, Tx_n))$ is nonincreasing so that $d(x_n, Tx_n) \rightarrow 0 \in \forall \exists$.

Applicability of Metatheorem

- Applied to asymptotic regularity statements $d(x_n, Tx_n) \rightarrow 0$, the corollary often gives full rates of convergence, e.g. because $(d(x_n, Tx_n))$ is nonincreasing so that $d(x_n, Tx_n) \rightarrow 0 \in \forall\exists$.
- From proofs of the convergence of (x_n) itself, one may only get **rates of metastability** Φ (Kreisel 1951, K.05, Tao 07) s.t.

$$\forall k \in \mathbb{N} \forall g \in \mathbb{N}^{\mathbb{N}} \exists n \in \mathbb{N} \forall i, j \in [n, n+g(n)] (d(x_i, x_j) < 2^{-k}) \in \forall\exists.$$

- **Admissible abstract structures:** metric, hyperbolic, $\text{CAT}(0)$, $\text{CAT}(\kappa > 0)$, Ptolemy, normed, their completions, Hilbert, uniformly convex, uniformly smooth (not: separable, strictly convex or smooth) spaces, abstract L^p - and $C(K)$ -spaces (and all other normed structures axiomatizable in positive bounded logic (in the sense of Henson, Iovino, Ben-Yaacov etc.).

- **Admissible abstract structures:** metric, hyperbolic, $\text{CAT}(0)$, $\text{CAT}(\kappa > 0)$, Ptolemy, normed, their completions, Hilbert, uniformly convex, uniformly smooth (not: separable, strictly convex or smooth) spaces, abstract L^p - and $C(K)$ -spaces (and all other normed structures axiomatizable in positive bounded logic (in the sense of Henson, Iovino, Ben-Yaacov etc.).
- **Admissible classes of functions:** uniformly continuous, Lipschitzian, nonexpansive, firmly- and strongly nonexpansive functions; also some classes of **discontinuous** functions: pseudo-contractions, maps with Suzuki's condition (E) etc.

- **Admissible abstract structures:** metric, hyperbolic, $CAT(0)$, $CAT(\kappa > 0)$, Ptolemy, normed, their completions, Hilbert, uniformly convex, uniformly smooth (not: separable, strictly convex or smooth) spaces, abstract L^p - and $C(K)$ -spaces (and all other normed structures axiomatizable in positive bounded logic (in the sense of Henson, Iovino, Ben-Yaacov etc.).
- **Admissible classes of functions:** uniformly continuous, Lipschitzian, nonexpansive, firmly- and strongly nonexpansive functions; also some classes of **discontinuous** functions: pseudo-contractions, maps with Suzuki's condition (E) etc.
Recently: set-valued accretive operators (Cauchy problems). (K./Koutsoukou-Argyraki, K./Powell).

- Uses of **ultraproducts** made in model theory can often be replaced by a **proof-theoretic uniform boundedness principle UB** which can be **eliminated** from proofs without contributing to the extracted bounds (K. ENTCS 2006, Engracia 2009, Günzel/K. Adv. Math. 2016). Recently **UB** has been used to replace sequential weak compactness (Ferreira, Leuştean, Pinto, Adv. Math. to appear).

Experience from numerous case studies

- Essentially, all **rates of asymptotic regularity** extracted have a very low complexity (**polynomial or exponential** in the data) at least in the most relevant special cases.

Experience from numerous case studies

- Essentially, all **rates of asymptotic regularity** extracted have a very low complexity (**polynomial or exponential** in the data) at least in the most relevant special cases.
- Except for 2 cases, all **rates of metastability** are of essentially the form

$$\Phi(\underline{a}, g) = (\chi_1(\underline{a}) \circ g \circ \chi_2(\underline{a}))^{B(\underline{a})}(0)$$

for simple (essentially polynomial) functions χ_1, χ_2, B in majorants \underline{a} of the parameters of the problem.

Experience from numerous case studies

- Essentially, all **rates of asymptotic regularity** extracted have a very low complexity (**polynomial or exponential** in the data) at least in the most relevant special cases.
- Except for 2 cases, all **rates of metastability** are of essentially the form

$$\Phi(\underline{a}, g) = (\chi_1(\underline{a}) \circ g \circ \chi_2(\underline{a}))^{B(\underline{a})}(0)$$

for simple (essentially polynomial) functions χ_1, χ_2, B in majorants \underline{a} of the parameters of the problem.

Implies: **algorithmic learnability** of a rate of convergence which - if a gap condition is satisfied - yields **oscillation bounds** (K./Safarik APAL 2014, Avigad/Rute ETDS 2015).

Proof-theoretic versus model-theoretic tameness

- In the recent book 'Model Theory and the Philosophy of Mathematical Practice: Formalization without Foundationalism', John Baldwin has argued that model theory became successful in applications to core mathematics by focusing on **local** foundations/formalizations rather than **global** ones and on **tame structures** (e.g. o-minimal ones).

Proof-theoretic versus model-theoretic tameness

- In the recent book 'Model Theory and the Philosophy of Mathematical Practice: Formalization without Foundationalism', John Baldwin has argued that model theory became successful in applications to core mathematics by focusing on **local** foundations/formalizations rather than **global** ones and on **tame structures** (e.g. o-minimal ones).
- We argue, that in a related way, also 'proof mining' is successful by focusing on **specific classes of problems** (e.g. iterations of nonlinear operators $T : C \rightarrow C$ on general convex subsets of abstract classes of normed or geodesic spaces).

Proof-theoretic versus model-theoretic tameness

- In contrast to model-theoretic tameness, quantification over \mathbb{N} and inductions etc. are crucially used in connection with convergence statements so that **Gödel-phenomena could occur in principle.**

Proof-theoretic versus model-theoretic tameness

- In contrast to model-theoretic tameness, quantification over \mathbb{N} and inductions etc. are crucially used in connection with convergence statements so that **Gödel-phenomena could occur in principle.**
- A different form of **'proof-theoretic tameness'** in existing ordinary (nonlinear) analysis largely leads **to extractable bounds of very low complexity.**

Proof-theoretic versus model-theoretic tameness

- In contrast to model-theoretic tameness, quantification over \mathbb{N} and inductions etc. are crucially used in connection with convergence statements so that **Gödel-phenomena could occur in principle.**
- A different form of **'proof-theoretic tameness'** in existing ordinary (nonlinear) analysis largely leads **to extractable bounds of very low complexity.**
- **Geometric properties** such as uniform convexity and smoothness etc. **more important than complicated inductions.**

Proof-theoretic versus model-theoretic tameness

- To **detect proof-theoretic tameness requires** to actually carry out the **proof analysis** (though usually some rough upper bound on the complexity can be obtained from proof-theoretic conservation results).

**Proof-theoretic tameness in practice I:
Polynomial rate of asymptotic regularity in
Bauschke's solution of the 'zero displacement
conjecture'**

Consider a Hilbert space H and nonempty closed and convex subsets $C_1, \dots, C_N \subseteq H$ with metric projections P_{C_i} , define $T := P_{C_N} \circ \dots \circ P_{C_1}$. In 2003 Bauschke proved the ‘zero displacement conjecture’:

$$\|T^{n+1}x - T^n x\| \rightarrow 0 \quad (x \in H).$$

Previously only known for $N = 2$ or $\text{Fix}(T) \neq \emptyset$ (or even $\bigcap_{i=1}^N C_i \neq \emptyset$) or C_i half spaces etc. starting with von Neumann.

Consider a Hilbert space H and nonempty closed and convex subsets $C_1, \dots, C_N \subseteq H$ with metric projections P_{C_i} , define $T := P_{C_N} \circ \dots \circ P_{C_1}$. In 2003 Bauschke proved the ‘zero displacement conjecture’:

$$\|T^{n+1}x - T^n x\| \rightarrow 0 \quad (x \in H).$$

Previously only known for $N = 2$ or $\text{Fix}(T) \neq \emptyset$ (or even $\bigcap_{i=1}^N C_i \neq \emptyset$) or C_i half spaces etc. starting with von Neumann. Proof uses abstract theory of maximal monotone operators: Minty’s theorem, Brézis-Haraux theorem, Rockafellar’s maximal monotonicity and sum theorems, strongly nonexpansive mappings, conjugate functions, normal cone operator...).

$(\|T^{n+1}x - T^n x\|)_{n \in \mathbb{N}}$ is **nonincreasing** and hence the conclusion in Bauschke's theorem is of the form $\forall \epsilon \exists$.

$(\|T^{n+1}x - T^n x\|)_{n \in \mathbb{N}}$ is **nonincreasing** and hence the conclusion in Bauschke's theorem is of the form $\forall \epsilon > 0, \exists N \in \mathbb{N}$.

Logical metatheorems, therefore, guarantee the extractability a uniform rate of asymptotic regularity which only depends on the error $\epsilon > 0$, $N \in \mathbb{N}$ and **majorants** for $x \in H$ and P_{C_1}, \dots, P_{C_N} : $b \geq \|x\|$ and $K \geq \|c_1\|, \dots, \|c_N\|$ for some **arbitrary** points $c_1 \in C_1, \dots, c_N \in C_N$:

$$\|P_{C_i} 0\| \leq \|c_i\| \leq K.$$

$(\|T^{n+1}x - T^n x\|)_{n \in \mathbb{N}}$ is **nonincreasing** and hence the conclusion in Bauschke's theorem is of the form $\forall \exists$.

Logical metatheorems, therefore, guarantee the extractability a uniform rate of asymptotic regularity which only depends on the error $\varepsilon > 0$, $N \in \mathbb{N}$ and **majorants** for $x \in H$ and P_{C_1}, \dots, P_{C_N} : $b \geq \|x\|$ and $K \geq \|c_1\|, \dots, \|c_N\|$ for some **arbitrary** points $c_1 \in C_1, \dots, c_N \in C_N$:

$$\|P_{C_i} 0\| \leq \|c_i\| \leq K.$$

Since the mappings P_{C_i} are nonexpansive, the corollary guarantees a computable $\Phi(\varepsilon, N, b, K)$ s.t. for $b \geq \|x\|$

$$\forall \varepsilon > 0 \forall n \geq \Phi(\varepsilon, N, b, K) (\|T^{n+1}x - T^n x\| < \varepsilon).$$

Theorem (K. FoCM 2019)

$$\Phi(\varepsilon, N, b, K) := \left\lceil \frac{18b + 12\alpha(\varepsilon/6)}{\varepsilon} - 1 \right\rceil \left\lceil \left(\frac{D}{\omega(D, \tilde{\varepsilon})} \right) \right\rceil$$

is a **rate of asymptotic regularity** in Bauschke's result, where

$$\tilde{\varepsilon} := \frac{\varepsilon^2}{27b + 18\alpha(\varepsilon/6)}, D := 2b + NK, \omega(D, \tilde{\varepsilon}) := \frac{1}{16D}(\tilde{\varepsilon}/N)^2.$$

$$\alpha(\varepsilon) := \frac{(K^2 + N^3(N-1)^2K^2)N^2}{\varepsilon}.$$

Here $b \geq \|x\|$ and $K \geq \left(\sum_{i=1}^N \|c_i\|^2 \right)^{\frac{1}{2}}$ for some $(c_1, \dots, c_N) \in C_1 \times \dots \times C_N$.

Proof-theoretic tameness in practice II:

Pursuit-evasion games: Lion-Man

Let (X, d) be a uniquely geodesic space, $D > 0$. $L_0, M_0 \in A$ starting points of the lion L and the man M . After n -steps, M moves to any point M_n s.t. $d(M_n, M_{n+1}) \leq D$ and L moves via the geodesic $[L_n, M_n]$ s.t. $d(L_n, L_{n+1}) = \min\{D, d(L_n, M_n)\}$.

Let (X, d) be a uniquely geodesic space, $D > 0$. $L_0, M_0 \in A$ starting points of the lion L and the man M . After n -steps, M moves to any point M_n s.t. $d(M_n, M_{n+1}) \leq D$ and L moves via the geodesic $[L_n, M_n]$ s.t. $d(L_n, L_{n+1}) = \min\{D, d(L_n, M_n)\}$.

López-Acedo/Nicolae/Piatek, Geom.Dedicat. to appear: if X is a compact uniquely geodesic space with the betweenness property, then **the lion wins** i.e. $\lim d(L_{n+1}, M_n) = 0$ (proof makes iterated use of sequential compactness, i.e. arithmetic comprehension **ACA**).

Let (X, d) be a uniquely geodesic space, $D > 0$. $L_0, M_0 \in A$ starting points of the lion L and the man M . After n -steps, M moves to any point M_n s.t. $d(M_n, M_{n+1}) \leq D$ and L moves via the geodesic $[L_n, M_n]$ s.t. $d(L_n, L_{n+1}) = \min\{D, d(L_n, M_n)\}$.

López-Acedo/Nicolae/Piatek, Geom.Dedicat. to appear: if X is a compact uniquely geodesic space with the betweenness property, then **the lion wins** i.e. $\lim d(L_{n+1}, M_n) = 0$ (proof makes iterated use of sequential compactness, i.e. arithmetic comprehension **ACA**).

' $\lim d(L_{n+1}, M_n) = 0$ ' $\in \Pi_2^0$ since the sequence is nonincreasing!

- **Proof Mining** extracts an explicit rate of convergence if one upgrades ‘uniquely geodesic’ and ‘betweenness property’ to ‘**uniform uniquely geodesic (with modulus)**’ and ‘**uniform betweenness property (with modulus Θ)**’.

- **Proof Mining** extracts an explicit rate of convergence if one upgrades ‘uniquely geodesic’ and ‘betweenness property’ to ‘**uniform uniquely geodesic (with modulus)**’ and ‘**uniform betweenness property (with modulus Θ)**’.
- With these upgrades the assumption of **compactness can be replaced by boundedness** in this particular case!

- **Proof Mining** extracts an **explicit rate of convergence** if one upgrades ‘uniquely geodesic’ and ‘betweenness property’ to ‘**uniform uniquely geodesic (with modulus)**’ and ‘**uniform betweenness property (with modulus Θ)**’.
- With these upgrades the assumption of **compactness can be replaced by boundedness** in this particular case!
- Even the **uniqueness of geodesics can be dropped**.

- **Proof Mining** extracts an **explicit rate of convergence** if one upgrades ‘uniquely geodesic’ and ‘betweenness property’ to ‘**uniform uniquely geodesic (with modulus)**’ and ‘**uniform betweenness property (with modulus Θ)**’.
- With these upgrades the assumption of **compactness can be replaced by boundedness** in this particular case!
- Even the **uniqueness of geodesics can be dropped**.
- Proof mining provides an explicit **rate of convergence** which only depends on Θ (in addition to $b \geq \text{diam}(A)$, $D, \varepsilon > 0$).

- **Proof Mining** extracts an **explicit rate of convergence** if one upgrades ‘uniquely geodesic’ and ‘betweenness property’ to ‘**uniform uniquely geodesic (with modulus)**’ and ‘**uniform betweenness property (with modulus Θ)**’.
- With these upgrades the assumption of **compactness can be replaced by boundedness** in this particular case!
- Even the **uniqueness of geodesics can be dropped**.
- Proof mining provides an explicit **rate of convergence** which only depends on Θ (in addition to $b \geq \text{diam}(A)$, $D, \varepsilon > 0$).
- **Moduli of uniform betweenness** can be **extracted** from **proofs of mere betweenness** for the admissible structures.

Betweenness and uniform betweenness in metric spaces

Definition (Diminnie and White 1981)

Let (X, d) be a metric space. X satisfies the betweenness property if for any distinct points $x, y, z, w \in X$

$$\left. \begin{array}{l} d(x, y) + d(y, z) \leq d(x, z) \\ d(y, z) + d(z, w) \leq d(y, w) \end{array} \right\} \Rightarrow d(x, z) + d(z, w) \leq d(x, w).$$

Betweenness and uniform betweenness in metric spaces

Definition (Diminnie and White 1981)

Let (X, d) be a metric space. X satisfies the betweenness property if for any distinct points $x, y, z, w \in X$

$$\left. \begin{array}{l} d(x, y) + d(y, z) \leq d(x, z) \\ d(y, z) + d(z, w) \leq d(y, w) \end{array} \right\} \Rightarrow d(x, z) + d(z, w) \leq d(x, w).$$

For normed spaces, betweenness follows from (but is strictly weaker than) strict convexity. It fails for $(\mathbb{R}^2, \|\cdot\|_\infty)$, $(\mathbb{R}^2, \|\cdot\|_1)$ but holds for some nonstrictly convex spaces.

The functional interpretation upgrades betweenness to (equivalent in the compact case!):

Definition (K., López-Acedo, Nicolae 2019)

A metric space (X, d) satisfies the uniform betweenness property with modulus $\Theta : (0, \infty)^3 \rightarrow (0, \infty)$ if

$$\forall \varepsilon, a, b > 0 \forall x, y, z, w \in X$$

$$\left(\left\{ \begin{array}{l} \text{sep}\{x, y, z, w\} \geq a \wedge \text{diam}\{x, y, z, w\} \leq b \\ d(x, y) + d(y, z) \leq d(x, z) + \Theta(\varepsilon, a, b) \\ d(y, z) + d(z, w) \leq d(y, w) + \Theta(\varepsilon, a, b) \\ \Rightarrow d(x, z) + d(z, w) \leq d(x, w) + \varepsilon \end{array} \right\} \right).$$

Definition (Lion-Man Game in general metric spaces)

X metric space, $D > 0$, $(M_n), (L_n)$ be sequences in X s.t.

$$d(M_n, M_{n+1}) \leq D, \quad d(L_{n+1}, L_n) + d(L_{n+1}, M_n) = d(L_n, M_n),$$

$$d(L_n, L_{n+1}) = \min\{D, d(L_n, M_n)\}.$$

Definition (Lion-Man Game in general metric spaces)

X metric space, $D > 0$, $(M_n), (L_n)$ be sequences in X s.t.

$$d(M_n, M_{n+1}) \leq D, \quad d(L_{n+1}, L_n) + d(L_{n+1}, M_n) = d(L_n, M_n),$$

$$d(L_n, L_{n+1}) = \min\{D, d(L_n, M_n)\}.$$

Then $\langle (M_n), (L_n) \rangle$ is a **Lion-Man game** with speed $D > 0$.

Definition (Lion-Man Game in general metric spaces)

\mathbf{X} metric space, $D > 0$, $(M_n), (L_n)$ be sequences in \mathbf{X} s.t.

$$d(M_n, M_{n+1}) \leq D, \quad d(L_{n+1}, L_n) + d(L_{n+1}, M_n) = d(L_n, M_n), \\ d(L_n, L_{n+1}) = \min\{D, d(L_n, M_n)\}.$$

Then $\langle (M_n), (L_n) \rangle$ is a **Lion-Man game** with speed $D > 0$.

Let \mathbf{X} be a **b -bounded** metric space with the uniform betweenness property with modulus Θ satisfying

$$\Theta(\varepsilon) := \Theta(\varepsilon, \varepsilon, b) \leq \varepsilon \quad \text{for all } \varepsilon > 0.$$

For $D > 0$ let $N \in \mathbb{N}$ be s.t. $b + 1 < ND$.

Theorem (K./López-Acedo/Nicolae 2019)

Let X be a bounded metric space with the uniform betweenness property and $\langle (M_n), (L_n) \rangle$ be a Lion-Man game, speed $D > 0$. Then the Lion approaches the man arbitrarily close.

Theorem (K./Lopéz-Acedo/Nicolae 2019)

Let X be a bounded metric space with the uniform betweenness property and $\langle (M_n), (L_n) \rangle$ be a Lion-Man game, speed $D > 0$. Then the Lion approaches the man arbitrarily close.

Moreover with $b \geq \text{diam}(X)$, Θ , N as above:

$$\forall \varepsilon > 0 \forall n \geq \Omega_{D,b,\Theta}(\varepsilon) \ (d(L_{n+1}, M_n) < \varepsilon),$$

where

$$\Omega_{D,b,\Theta}(\varepsilon) = N + N \left\lceil \frac{b}{\Theta^{(N)}(\alpha)} \right\rceil$$

with

$$0 < \alpha \leq \min \left\{ \frac{1}{N}, \frac{D}{2}, \frac{\varepsilon}{2} \right\}.$$

Moduli of uniform betweenness

Θ can be **explicitly computed** for L^p ($1 < p < \infty$) and $\text{CAT}(\kappa)$ -spaces, $\kappa > 0$.

Moduli of uniform betweenness

Θ can be **explicitly computed** for L^p ($1 < p < \infty$) and $\text{CAT}(\kappa)$ -spaces, $\kappa > 0$.

$$\Theta(\varepsilon, a, b) = \min \left\{ \Phi \left(\min \left\{ \frac{a \cdot \varepsilon}{8b}, \frac{a}{2} \right\}, b \right), a \right\} \quad \text{with}$$

Moduli of uniform betweenness

Θ can be **explicitly computed** for L^p ($1 < p < \infty$) and $\text{CAT}(\kappa)$ -spaces, $\kappa > 0$.

$$\Theta(\varepsilon, a, b) = \min \left\{ \Phi \left(\min \left\{ \frac{a \cdot \varepsilon}{8b}, \frac{a}{2} \right\}, b \right), a \right\} \quad \text{with}$$

For L^p :

$$\Phi(\varepsilon, b) = \begin{cases} \frac{p-1}{8} \frac{\varepsilon^2}{(b+\varepsilon)}, & \text{if } 1 < p \leq 2, \\ \frac{1}{p2^p} \frac{\varepsilon^p}{(b+\varepsilon)^{p-1}}, & \text{if } 2 < p < \infty. \end{cases}$$

Moduli of uniform betweenness

Θ can be **explicitly computed** for L^p ($1 < p < \infty$) and $\text{CAT}(\kappa)$ -spaces, $\kappa > 0$.

$$\Theta(\varepsilon, a, b) = \min \left\{ \Phi \left(\min \left\{ \frac{a \cdot \varepsilon}{8b}, \frac{a}{2} \right\}, b \right), a \right\} \quad \text{with}$$

For L^p :

$$\Phi(\varepsilon, b) = \begin{cases} \frac{p-1}{8} \frac{\varepsilon^2}{(b+\varepsilon)}, & \text{if } 1 < p \leq 2, \\ \frac{1}{p2^p} \frac{\varepsilon^p}{(b+\varepsilon)^{p-1}}, & \text{if } 2 < p < \infty. \end{cases}$$

For $\text{CAT}(\kappa)$ -spaces X , $\kappa > 0$, with $\text{diam}(X) < \pi/(2\sqrt{\kappa})$:

$$\Phi(\varepsilon, b) = \frac{c}{16} \frac{\varepsilon^2}{b + \varepsilon}, \quad \text{where}$$

$c = (\pi - 2\sqrt{\kappa} \beta) \tan(\sqrt{\kappa} \beta)$ for any $0 < \beta \leq \pi/(2\sqrt{\kappa}) - \text{diam}(X)$.

Ptolemy spaces

Definition

A metric space (X, d) is **Ptolemy** if for all $x, y, z, w \in X$

$$d(x, z)d(y, w) \leq d(x, y)d(z, w) + d(x, w)d(y, z).$$

Ptolemy spaces

Definition

A metric space (X, d) is **Ptolemy** if for all $x, y, z, w \in X$

$$d(x, z)d(y, w) \leq d(x, y)d(z, w) + d(x, w)d(y, z).$$

Proposition (Foertsch, Lytchak, Schroeder 2007)

There are complete bounded Ptolemy spaces which are geodesic but **not uniquely geodesic**.

Ptolemy spaces

Definition

A metric space (X, d) is **Ptolemy** if for all $x, y, z, w \in X$

$$d(x, z)d(y, w) \leq d(x, y)d(z, w) + d(x, w)d(y, z).$$

Proposition (Foertsch, Lytchak, Schroeder 2007)

There are complete bounded Ptolemy spaces which are geodesic but **not uniquely geodesic**.

Proposition (K., López-Acedo, Nicolae 2019)

Let (X, d) be a Ptolemy space. Then

$\Theta(\varepsilon, a, b) := \sqrt{b^2 + \varepsilon a} - b$ is a modulus for the uniform betweenness property.

A borderline case for proof-theoretic tameness

U. Kohlenbach, A. Sipoş, The finitary content of sunny nonexpansive retractions. arXiv:1812.04940 [math.FA], 2018.

The Browder-Halpern result

Let $C \subseteq H$ be a bounded, closed and convex subset of a Hilbert space H . $T : C \rightarrow C$ be nonexpansive, $x_0 \in C$ and $t \in [0, 1)$.

$$T_t : C \rightarrow C, \quad T_t(x) := tTx + (1 - t)x_0$$

is a t -contraction and so has a unique point x_t with $x_t = T_t x_t$.

The Browder-Halpern result

Let $C \subseteq H$ be a bounded, closed and convex subset of a Hilbert space H . $T : C \rightarrow C$ be nonexpansive, $x_0 \in C$ and $t \in [0, 1)$.

$$T_t : C \rightarrow C, \quad T_t(x) := tTx + (1 - t)x_0$$

is a t -contraction and so has a unique point x_t with $x_t = T_t x_t$.

Theorem (Browder 1967; Halpern 1967)

$p := \lim_{t \rightarrow 1} x_t$ exists strongly and $p = P_{\text{Fix}(T)}x$, where P denotes the metric projection.

The Browder-Halpern result

Let $C \subseteq H$ be a bounded, closed and convex subset of a Hilbert space H . $T : C \rightarrow C$ be nonexpansive, $x_0 \in C$ and $t \in [0, 1)$.

$$T_t : C \rightarrow C, \quad T_t(x) := tTx + (1 - t)x_0$$

is a t -contraction and so has a unique point x_t with $x_t = T_t x_t$.

Theorem (Browder 1967; Halpern 1967)

$p := \lim_{t \rightarrow 1} x_t$ exists strongly and $p = P_{\text{Fix}(T)}x$, where P denotes the metric projection.

Even in simple cases on $[0, 1]$ there is in general **no computable rate convergence**. However, a **primitive recursive in the simple form as mentioned above rate of metastability** is extracted in (K., Adv. Math. 2011).

S. Reich's Theorem

Even for L^p ($1 < p < \infty, p \neq 2$) open until the celebrated:

S. Reich's Theorem

Even for L^p ($1 < p < \infty, p \neq 2$) open until the celebrated:

Theorem (S. Reich, 1980)

In the framework above, if X is a **uniformly smooth Banach space**, then for all $x \in C$ we have that $\lim_{t \rightarrow 1} x_t := p$ exists and it is a fixed point of T . Moreover $p = Q_{\text{Fix}(T)}x$, where Q is the unique **sunny nonexpansive retraction** onto $\text{Fix}(T)$.

S. Reich's Theorem

Even for L^p ($1 < p < \infty, p \neq 2$) open until the celebrated:

Theorem (S. Reich, 1980)

In the framework above, if X is a **uniformly smooth Banach space**, then for all $x \in C$ we have that $\lim_{t \rightarrow 1} x_t := p$ exists and it is a fixed point of T . Moreover $p = Q_{\text{Fix}(T)}x$, where Q is the unique **sunny nonexpansive retraction** onto $\text{Fix}(T)$.

$Q = P$ iff X is a Hilbert space (Bruck 1974).

S. Reich's Theorem

Even for L^p ($1 < p < \infty, p \neq 2$) open until the celebrated:

Theorem (S. Reich, 1980)

In the framework above, if X is a **uniformly smooth Banach space**, then for all $x \in C$ we have that $\lim_{t \rightarrow 1} x_t := p$ exists and it is a fixed point of T . Moreover $p = Q_{\text{Fix}(T)}x$, where Q is the unique **sunny nonexpansive retraction** onto $\text{Fix}(T)$.

$Q = P$ iff X is a Hilbert space (Bruck 1974).

The convergence of numerous iterative algorithms in nonlinear analysis is based on Reich's theorem!

Sunny nonexpansive retractions

Let E be a nonempty subset of C and $Q : C \rightarrow E$. We call Q a **retraction** if for all $x \in E$, $Qx = x$. If Q is a retraction, we call it **sunny** if for all $x \in C$ and $t \geq 0$, $Q(Qx + t(x - Qx)) = Qx$.

Sunny nonexpansive retractions

Let E be a nonempty subset of C and $Q : C \rightarrow E$. We call Q a **retraction** if for all $x \in E$, $Qx = x$. If Q is a retraction, we call it **sunny** if for all $x \in C$ and $t \geq 0$, $Q(Qx + t(x - Qx)) = Qx$.

Proposition (Variational Inequality)

A retraction $Q : C \rightarrow E$ is sunny and nonexpansive iff for all $x \in C$ and $y \in E$,

$$\langle x - Qx, j(y - Qx) \rangle \leq 0.$$

Sunny nonexpansive retractions

Let E be a nonempty subset of C and $Q : C \rightarrow E$. We call Q a **retraction** if for all $x \in E$, $Qx = x$. If Q is a retraction, we call it **sunny** if for all $x \in C$ and $t \geq 0$, $Q(Qx + t(x - Qx)) = Qx$.

Proposition (Variational Inequality)

A retraction $Q : C \rightarrow E$ is sunny and nonexpansive iff for all $x \in C$ and $y \in E$,

$$\langle x - Qx, j(y - Qx) \rangle \leq 0.$$

As a consequence, there is at most one sunny nonexpansive retraction $Q : C \rightarrow E$ (Bruck 1973).

Sunny nonexpansive retractions

Let E be a nonempty subset of C and $Q : C \rightarrow E$. We call Q a **retraction** if for all $x \in E$, $Qx = x$. If Q is a retraction, we call it **sunny** if for all $x \in C$ and $t \geq 0$, $Q(Qx + t(x - Qx)) = Qx$.

Proposition (Variational Inequality)

A retraction $Q : C \rightarrow E$ is sunny and nonexpansive iff for all $x \in C$ and $y \in E$,

$$\langle x - Qx, j(y - Qx) \rangle \leq 0.$$

As a consequence, there is at most one sunny nonexpansive retraction $Q : C \rightarrow E$ (Bruck 1973).

The existence of (sunny) nonexpansive retractions onto $\text{Fix}(T)$ was first shown by R. Bruck in 1971, 1973 using Zorn's lemma.

One key step in the proof

Consider $f : C \rightarrow \mathbb{R}_+$ with $f(z) := \limsup_{n \rightarrow \infty} \|x_n - z\|$. Let K be the set of minimizers of f . Claim: $K \cap \text{Fix}(T) \neq \emptyset$.

One key step in the proof

Consider $f : C \rightarrow \mathbb{R}_+$ with $f(z) := \limsup_{n \rightarrow \infty} \|x_n - z\|$. Let K be the set of minimizers of f . Claim: $K \cap \text{Fix}(T) \neq \emptyset$.

Since f is convex and continuous, C is closed convex bounded nonempty, and X is uniformly smooth, hence reflexive, we have that $K \neq \emptyset$.

One key step in the proof

Consider $f : C \rightarrow \mathbb{R}_+$ with $f(z) := \limsup_{n \rightarrow \infty} \|x_n - z\|$. Let K be the set of minimizers of f . Claim: $K \cap \text{Fix}(T) \neq \emptyset$.

Since f is convex and continuous, C is closed convex bounded nonempty, and X is uniformly smooth, hence reflexive, we have that $K \neq \emptyset$. Let $y \in K$ and $z \in C$. Then:

$$\begin{aligned} f(Ty) &= \limsup_{n \rightarrow \infty} \|x_n - Ty\| \leq \limsup_{n \rightarrow \infty} (\|x_n - Tx_n\| + \|Tx_n - Ty\|) \\ &\leq \limsup_{n \rightarrow \infty} (\|x_n - Tx_n\| + \|x_n - y\|) \\ &\leq \limsup_{n \rightarrow \infty} \|x_n - Tx_n\| + \limsup_{n \rightarrow \infty} \|x_n - y\| \\ &= f(y) \leq f(z), \end{aligned}$$

so $Ty \in K$.

One key step in the proof

Consider $f : C \rightarrow \mathbb{R}_+$ with $f(z) := \limsup_{n \rightarrow \infty} \|x_n - z\|$. Let K be the set of minimizers of f . Claim: $K \cap \text{Fix}(T) \neq \emptyset$.

Since f is convex and continuous, C is closed convex bounded nonempty, and X is uniformly smooth, hence reflexive, we have that $K \neq \emptyset$. Let $y \in K$ and $z \in C$. Then:

$$\begin{aligned} f(Ty) &= \limsup_{n \rightarrow \infty} \|x_n - Ty\| \leq \limsup_{n \rightarrow \infty} (\|x_n - Tx_n\| + \|Tx_n - Ty\|) \\ &\leq \limsup_{n \rightarrow \infty} (\|x_n - Tx_n\| + \|x_n - y\|) \\ &\leq \limsup_{n \rightarrow \infty} \|x_n - Tx_n\| + \limsup_{n \rightarrow \infty} \|x_n - y\| \\ &= f(y) \leq f(z), \end{aligned}$$

so $Ty \in K$. Since K is a closed convex bounded nonempty T -invariant subset of a uniformly smooth space, there is a $p \in K \cap \text{Fix}(T)$.

The proof gets somewhat easier if X is assumed to be also **uniformly convex** (still covering L^p -spaces): only **arbitrarily good ε -minimizers** needed.

The proof gets somewhat easier if X is assumed to be also **uniformly convex** (still covering L^p -spaces): only **arbitrarily good ε -minimizers** needed.

Next: replace **lim sup**'s (\Leftrightarrow ACA) by **approximate lim sup**'s ($\Leftrightarrow \Pi_2^0$ -IA), in a process known as **arithmetization** (K.1996).

The proof gets somewhat easier if X is assumed to be also **uniformly convex** (still covering L^p -spaces): only **arbitrarily good ε -minimizers** needed.

Next: replace **lim sup**'s (\Leftrightarrow ACA) by **approximate lim sup**'s ($\Leftrightarrow \Pi_2^0$ -IA), in a process known as **arithmetization** (K.1996).

While the functional interpretation of the ε -minimization in general requires Gödel primitive recursion of type 2 (**T_2**) the actual extraction constructs a **rate of metastability** in **$\Phi \in T_1$** (i.e. of Ackermann type).

The proof gets somewhat easier if X is assumed to be also **uniformly convex** (still covering L^p -spaces): only **arbitrarily good ε -minimizers** needed.

Next: replace **lim sup**'s (\Leftrightarrow ACA) by **approximate lim sup**'s ($\Leftrightarrow \Pi_2^0$ -IA), in a process known as **arithmetization** (K.1996).

While the functional interpretation of the ε -minimization in general requires Gödel primitive recursion of type 2 (**T_2**) the actual extraction constructs a **rate of metastability** in **$\Phi \in T_1$** (i.e. of Ackermann type).

It may well be that a closer analysis of Φ shows that it is already definable in **T_0** (in line with a classical result of Parsons that certain forms of type-1 primitive recursion can be reduced to **T_0**).

Future of Proof Mining

- Find new ‘organic connections’ (Kreisel) between core mathematics and proof theory.

Future of Proof Mining

- Find new ‘organic connections’ (Kreisel) between core mathematics and proof theory.
- New classes of results rewarding for proof-theoretic analysis.

Future of Proof Mining

- Find new ‘organic connections’ (Kreisel) between core mathematics and proof theory.
- New classes of results rewarding for proof-theoretic analysis.
- Proofs which use highly abstract ‘ideal’ principles to prove concrete numerically meaningful results are most promising.

Future of Proof Mining

- Find new ‘organic connections’ (Kreisel) between core mathematics and proof theory.
- New classes of results rewarding for proof-theoretic analysis.
- Proofs which use highly abstract ‘ideal’ principles to prove concrete numerically meaningful results are most promising.
- Built suitable local proof-theoretic methods to cover such classes of proofs appropriately.

Future of Proof Mining

- Find new ‘organic connections’ (Kreisel) between core mathematics and proof theory.
- New classes of results rewarding for proof-theoretic analysis.
- Proofs which use highly abstract ‘ideal’ principles to prove concrete numerically meaningful results are most promising.
- Built suitable local proof-theoretic methods to cover such classes of proofs appropriately.
- The area of analysis has been particularly fruitful. But other promising areas: geometry, algebra (see Simmons/Towsner Adv.Math.).

Recent Surveys:

Recent Surveys:

U. Kohlenbach, Proof-Theoretic Methods in Nonlinear Analysis. In. Proc. ICM 2018, Proc. ICM 2018, B. Sirakov, P. Ney de Souza, M. Viana (eds.), Vol. 2, pp. 61-82. World Scientific 2019.

Recent Surveys:

U. Kohlenbach, Proof-Theoretic Methods in Nonlinear Analysis. In. Proc. ICM 2018, Proc. ICM 2018, B. Sirakov, P. Ney de Souza, M. Viana (eds.), Vol. 2, pp. 61-82. World Scientific 2019.

U. Kohlenbach, Local Formalizations in Nonlinear Analysis and Related Areas and Proof-Theoretic Tameness. To appear in forthcoming volume (eds. P. Weingartner, H.-P. Leeb) 'Kreisel's Interests - On the Foundations of Logic and Mathematics'.